

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

IDH2 (R172K)

Catalog Number: 26163

Gene Symbol: IDH2; D2HGA2; ICD-M; IDH; IDHM; IDP; IDPM; mNADP-IDH

Description: Anti-IDH2 (R172K) Mouse Monoclonal Antibody

Background: Isocitrate dehydrogenase (IDH) catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. The isocitrate and isopropylmalate dehydrogenases family has three members, IDH1, IDH2 and IDH3. IDH2 plays a role in intermediary metabolism and energy production. Defects in IDH2 are the cause of D-2-hydroxyglutaric aciduria type 2 (D2HGA2). Somatic mosaic mutations of this protein have also been found associated to Ollier disease and Maffucci syndrome, and R172K IDH2 mutations do exist in diffusely infiltrative gliomas.

Immunogen: A synthetic peptide from theinternal region of IDH2, which includes themutation of R172K, human origin.

Tested Applications: ELISA, WB, IF, IHC

Recommended Dilutions:

ELISA: 1:1000-1:5000 WB: 1:500-1:1000 IF: 1:50-1:100 IHC: 1:10-1:100

Concentration: 1 mg/ml

Host: Mouse

Clonality: Monoclonal

Isotype: IgG

Purity: Purified from ascites

Format: Liquid
Preservative: No

Constituents: PBS (without Mg²⁺ and Ca²⁺), pH7.4, 150

mM NaCl, 50% glycerol


Species Reactivity: Recognizes IDH2(R172K) mutant,

but not wild-type IDH2 of vertebrates.

Storage Conditions: Store at -20°C. Avoid freeze / thaw

cycles.

Western blot:



WB: anti-IDH2(R172K) mAb

Western blot analysis of recombinant IDH2(R172K) and wild type proteins.

Purified His-tagged IDH2 (R172K) (lane 2) and wildtypeprotein (lane 1) were blotted with anti-IDH2 (R172K) monoclonal antibody (Cat. # 26163).

Support:

Web:

610-945-2007


info@neweastbio.com

www.neweastbio.com



Pioneering GTPase and Oncogene Product Development since 2010

Immunofluorescence:



Immunofluorescence of cells expressing IDH2 proteins with anti-IDH2 (R172K) antibody.

HEK293T cells were transfected with pCDNA3-GFP- IDH2 (R172K) plasmid, pCDNA3-GFP- IDH2(WT) plasmid, pCDNA3-GFP- IDH2 (R172H) plasmid, pCDNA3-GFP- IDH2 (R172W) plasmid, pCDNA3-GFP- IDH2 (R172G) plasmid, pCDNA3-GFP- IDH2 (R172M) plasmid or pCDNA3-GFP- IDH2 (R172S) plasmid, then fixed and stained with anti- IDH2 (R172K) monoclonal antibody (Cat. #26163)

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS