

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

ACTC1 RABBIT PAB

Cat.#: S218667

Product Name: Anti-ACTC1 Rabbit Polyclonal Antibody

Synonyms: ACTC; ASD5; CMD1R; CMH11; LVNC4

UNIPROT ID: P68032 (Gene Accession - BC009978)

Background: Actins are highly conserved proteins that are involved in various types of cell motility. Polymerization of globular actin (G-actin) leads to a structural filament (F-actin) in the form of a two-stranded helix. Each actin can bind to four others. The protein encoded by this gene belongs to the actin family which is comprised of three main groups of actin isoforms, alpha, beta, and gamma. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. Defects in this gene have been associated with idiopathic dilated cardiomyopathy (IDC) and familial hypertrophic cardiomyopathy (FHC).

Immunogen: Fusion protein of human ACTC1

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-100;WB: 1000-5000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


Research Areas: Signal Transduction, Cardiovascular

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human gastric cancer tissue using 218667(ACTC1 Antibody) at a dilution of 1/50(Cytoplasm).

In comparision with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with the fusion protein and then with 218667(Anti-ACTC1 Antibody) at dilution 1/50.

kDa	
130—	
95 —	
72 —	
55—	
	-
36—	
36— 28—	

Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Mouse heart tissue lysate; Primary antibody: 218667(ACTC1 Antibody) at dilution 1/600; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 1 second