

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## AK4 RABBIT PAB

Cat.#: S221336

**Product Name:** Anti-AK4 Rabbit Polyclonal Antibody

Synonyms: AK3; AK 4; AK3L1; AK3L2

UNIPROT ID: P27144 (Gene Accession - NP\_037542)

**Background:** This gene encodes a member of the adenylate kinase family of enzymes. The encoded protein is localized to the mitochondrial matrix. Adenylate kinases regulate the adenine and guanine nucleotide compositions within a cell by catalyzing the reversible transfer of phosphate group among these nucleotides. Five isozymes of adenylate kinase have been identified in vertebrates. Expression of these isozymes is tissue–specific and developmentally regulated. A pseudogene for this gene has been located on chromosome 17. Three transcript variants encoding the same protein have been identified for this gene. Sequence alignment suggests that the gene defined by NM\_013410, NM\_203464, and NM\_001005353 is located on chromosome 1.

**Immunogen:** Synthetic peptide of human AK4

**Applications:** ELISA, WB, IHC

**Recommended Dilutions:** IHC: 25-100;WB: 200-1000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

**Isotype:** Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Metabolism

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human brain tissue using 221336(AK4 Antibody) at a dilution of 1/30(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human brain tissue is first treated with the synthetic peptide and then with 221336 (Anti-AK4 Antibody) at dilution 1/30.



Gel: 12%SDS-PAGE, Lysate: 40 µg;

Lane 1-2: MCF-7 cell and mouse kidney tissue lysates;

Primary antibody: 221336(AK4 Antibody) at

dilution 1/300;

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution;

Exposure time: 15 seconds