

## AOAH RABBIT PAB

**Cat.#:** S218757

**Product Name:** Anti-AOAH Rabbit Polyclonal Antibody

**Synonyms:**

**UNIPROT ID:** P28039 (Gene Accession - BC025698 )

**Background:** This locus encodes both the light and heavy subunits of acyloxyacyl hydrolase. The encoded enzyme catalyzes the hydrolysis of acyloxyacyl-linked fatty acyl chains from bacterial lipopolysaccharides, effectively detoxifying these molecules. The encoded protein may play a role in modulating host inflammatory response to gram-negative bacteria. Alternatively spliced transcript variants have been described.

**Immunogen:** Fusion protein of human AOAH

**Applications:** ELISA, WB, IHC

**Recommended Dilutions:** IHC: 20-100;WB: 200-1000;ELISA: 5000-10000

**Host Species:** Rabbit

**Clonality:** Rabbit Polyclonal

**Isotype:** Immunogen-specific rabbit IgG


**Purification:** Antigen affinity purification

**Species Reactivity:** Human, Mouse


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

**Research Areas:** Metabolism, Immunology


**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 218757(AOAH Antibody) at a dilution of 1/20(Secreted).



In comparison with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the fusion protein and then with 218757(Anti-AOAH Antibody) at dilution 1/20.



Gel: 8%SDS-PAGE, Lysate: 40 µg;  
Lane: HL-60 cell lysate;  
Primary antibody: 218757(AOAH Antibody) at dilution 1/250;  
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;  
Exposure time: 5 seconds