

## ATP5G RABBIT MAB

**Cat.#:** N261894

**Product Name:** Anti-ATP5G Rabbit Monoclonal Antibody

**Synonyms:** ATP synthase lipid-binding protein; ATP synthase membrane subunit c locus 1

**UNIPROT ID:** P05496/Q06055/P48201

**Background:** Mitochondrial membrane ATP synthase (F<sub>1</sub>F<sub>0</sub> ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F<sub>1</sub> - containing the extramembraneous catalytic core and F<sub>0</sub> - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F<sub>1</sub> is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F<sub>0</sub> domain. A homomeric c-ring of probably 10 subunits is part of the complex rotary element. Miscellaneous There are three genes which encode the mitochondrial ATP synthase proteolipid and they specify precursors with different import sequences but identical mature proteins. Is the major protein stored in the storage bodies of animals or humans affected with ceroid lipofuscinosis (Batten disease).

**Immunogen:** A synthetic peptide of human ATP5G1/G2/G3

**Applications:** WB, IHC-P, IP

**Recommended Dilutions:** WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20

**Host Species:** Rabbit

**Clonality:** Rabbit Monoclonal

**Clone ID:** R02-6G1

**MW:** Calculated MW: 14 kDa; Observed MW: 14 kDa

**Isotype:** IgG

**Purification:** Affinity Purified

**Species Reactivity:** Human, Rat


**Conjugation:** Unconjugated

**Modification:** Unmodified


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**Research Areas:** Signal Transduction

**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin-embedded Human tonsil using ATP5G1/G2/G3 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of ATP5G1/G2/G3 in Jurkat, C6 lysates using ATP5G antibody.