

ATP6V0D1 RABBIT MAB

Cat.#: N261895

Product Name: Anti-ATP6V0D1 Rabbit Monoclonal Antibody

Synonyms: P39; VATX; VMA6; ATP6D; ATP6DV; VPATPD

UNIPROT ID: P61421

Background: Subunit of the integral membrane V0 complex of vacuolar ATPase. Vacuolar ATPase is responsible for acidifying a variety of intracellular compartments in eukaryotic cells, thus providing most of the energy required for transport processes in the vacuolar system. May play a role in coupling of proton transport and ATP hydrolysis. May play a role in cilium biogenesis through regulation of the transport and the localization of proteins to the cilium. In aerobic conditions, involved in intracellular iron homeostasis, thus triggering the activity of Fe²⁺ prolyl hydroxylase (PHD) enzymes, and leading to HIF1A hydroxylation and subsequent proteasomal degradation (PubMed:28296633).

Immunogen: Recombinant protein of human ATP6V0D1

Applications: WB,IHC-F,IHC-P,ICC/IF,IP

Recommended Dilutions: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20

Host Species: Rabbit

Clonality: Rabbit Monoclonal

Clone ID: R07-1A5

MW: Calculated MW: 40 kDa; Observed MW: 40 kDa

Isotype: IgG

Purification: Affinity Purified

Species Reactivity: Human,Mouse,Rat


Conjugation: Unconjugated

Modification: Unmodified


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

Research Areas: Signal Transduction


Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Immunocytochemistry analysis of ATP6V0D1 (green) in HeLa using ATP6V0D1 antibody, and DAPI (blue)



Western blot analysis of ATP6V0D1 in K562, rat Brain, C6, 3T3, HeLa lysates using ATP6V0D1 antibody.



Immunohistochemistry analysis of paraffin-embedded Human colon cancer using ATP6V0D1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.