

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CCNG2 RABBIT PAB

Cat.#: S221262

Product Name: Anti-CCNG2 Rabbit Polyclonal Antibody

Synonyms:

UNIPROT ID: Q16589 (Gene Accession - NP_004345)

Background: The eukaryotic cell cycle is governed by cyclin-dependent protein kinases (CDKs) whose activities are regulated by cyclins and CDK inhibitors. The 8 species of cyclins reported in mammals, cyclins A through H, share a conserved amino acid sequence of about 90 residues called the cyclin box. The amino acid sequence of cyclin G is well conserved among mammals. The nucleotide sequence of cyclin G1 and cyclin G2 are 53% identical. Unlike cyclin G1, cyclin G2 contains a C-terminal PEST protein destabilization motif, suggesting that cyclin G2 expression is tightly regulated through the cell cycle.

Immunogen: Synthetic peptide of human CCNG2

Applications: ELISA, IHC

Recommended Dilutions: IHC: 25-100; ELISA: 2000-5000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Epigenetics and Nuclear Signaling, Cancer

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 221262(CCNG2 Antibody) at a dilution of 1/25(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 221262(Anti-CCNG2 Antibody) at dilution 1/25.



The image on the left is immunohistochemistry of paraffinembedded Human esophagus cancer tissue using 221262(Anti-CCNG2 Antibody) at a dilution of 1/25.



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with synthetic peptide and then with D262778 (Anti-CCNG2 Antibody) at dilution 1/25.