

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CD22 RABBIT PAB

Cat.#: S217260

Product Name: Anti-CD22 Rabbit Polyclonal Antibody

Synonyms: SIGLEC2; SIGLEC-2

UNIPROT ID: P20273 (Gene Accession - BC109306)

Background: The B lymphocyte specific CD22 antigen, also designated B-lymphocyte cell adhesion molecule (BLCAM), sialic acid-binding Ig-like lectin 2 (Siglec-2) and Leu-14, is a type I integral membrane glycoprotein, structurally similar to other cell adhesion molecules (CAMs), which acts as a regulator of B cell signaling. CD22 is expressed as both a cytoplasmic and membrane protein during discrete stages of B cell lymphocyte differentiation. The cytoplasmic form of CD22, expressed early in B cell development, is a useful marker for acute lymphocytic leukemia. The membrane form of CD22 is expressed in mature B cells prior to their differentiation into plasma cells.

Immunogen: Fusion protein of human CD22

Applications: ELISA, IHC

Recommended Dilutions: IHC: 50-200; ELISA: 2000-5000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification

Species Reactivity: Human

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Cancer, Immunology, Stem Cells

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human brain tissue using 217260(CD22 Antibody) at a dilution of 1/50(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human brain tissue is first treated with the fusion protein and then with 217260 (Anti-CD22 Antibody) at dilution 1/50.



The image on the left is immunohistochemistry of paraffinembedded Human thyroid cancer tissue using 217260(Anti-CD22 Antibody) at a dilution of 1/50.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with fusion protein and then with D222064(Anti-CD22 Antibody) at dilution 1/50.