

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CD24 RABBIT PAB

Cat.#: S220432

Product Name: Anti-CD24 Rabbit Polyclonal Antibody

Synonyms: CD24A

UNIPROT ID: P25063 (Gene Accession - NP_037362)

Background: This gene encodes a sialoglycoprotein that is expressed on mature granulocytes and B cells and modulates growth and differentiation signals to these cells. The precursor protein is cleaved to a short 32 amino acid mature peptide which is anchored via a glycosyl phosphatidylinositol (GPI) link to the cell surface. This gene was missing from previous genome assemblies, but is properly located on chromosome 6. Non-transcribed pseudogenes have been designated on chromosomes 1, 15, 20, and Y. Alternative splicing results in multiple transcript variants.

Immunogen: Synthetic peptide of human CD24

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-100;WB: 200-1000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human


Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Cancer, Cardiovascular, Metabolism, Neuroscience, Immunology, Stem Cells **Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human colorectal cancer tissue using 220432(CD24 Antibody) at a dilution of 1/50(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human colorectal cancer tissue is first treated with the synthetic peptide and then with 220432(Anti-CD24 Antibody) at dilution 1/50.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-2: Raji and LOVO cell lysates; Primary antibody: 220432(CD24 Antibody) at dilution 1/400; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 5 minutes