

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

CXCR4 RABBIT PAB

Cat.#: S221197

Product Name: Anti-CXCR4 Rabbit Polyclonal Antibody

Synonyms: FB22; HM89; LAP3; LCR1; NPYR; WHIM; CD184; LAP-3; LESTR; NPY3R; NPYRL; WHIMS; HSY3RR;

NPYY3R; D2S201E

UNIPROT ID: P61073 (Gene Accession - NP_003458)

Background: This gene encodes a CXC chemokine receptor specific for stromal cell-derived factor-1. The protein has 7 transmembrane regions and is located on the cell surface. It acts with the CD4 protein to support HIV entry into cells and is also highly expressed in breast cancer cells. Mutations in this gene have been associated with WHIM (warts, hypogammaglobulinemia, infections, and myelokathexis) syndrome. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Immunogen: Synthetic peptide of human CXCR4

Applications: ELISA, IHC

Recommended Dilutions: IHC: 25-100; ELISA: 2000-5000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Cancer, Cardiovascular, Metabolism, Neuroscience, Immunology, Stem Cells

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 221197(CXCR4 Antibody) at a dilution of 1/20(Cell membrane).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 221197(Anti-CXCR4 Antibody) at dilution 1/20.



The image on the left is immunohistochemistry of paraffinembedded Human esophagus cancer tissue using 221197(Anti-CXCR4 Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with synthetic peptide and then with D262693(Anti-CXCR4 Antibody) at dilution 1/20.