

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **CYCS RABBIT PAB**

**Cat.#:** S216463

**Product Name:** Anti-CYCS Rabbit Polyclonal Antibody

Synonyms: CYC; HCS; THC4

UNIPROT ID: P99999 (Gene Accession - BC015130)

**Background:** This gene encodes a small heme protein that functions as a central component of the electron transport chain in mitochondria. The encoded protein associates with the inner membrane of the mitochondrion where it accepts electrons from cytochrome b and transfers them to the cytochrome oxidase complex. This protein is also involved in initiation of apoptosis. Mutations in this gene are associated with autosomal dominant nonsyndromic thrombocytopenia.

Numerous processed pseudogenes of this gene are found throughout the human genome.

Immunogen: Fusion protein of human CYCS

**Applications:** ELISA, WB, IHC

**Recommended Dilutions:** IHC: 100-300;WB: 500-2000;ELISA: 2000-10000

Host Species: Rabbit

**Clonality:** Rabbit Polyclonal

**Isotype:** Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


**Research Areas:** Metabolism, Cancer, Cardiovascular

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffinembedded Human colon cancer tissue using 216463(CYCS Antibody) at a dilution of 1/70(Cytoplasm).



Immunohistochemistry analysis of paraffinembedded Human gastric cancer tissue using 216463(Anti-CYCS Antibody) at a dilution of 1/70.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-7: Hela cells, K562 cells, mouse brain tissue, Mouse heart tissue, Mouse kidney tissue, PC3 cells, Jurkat cells; Primary antibody: 216463(CYCS Antibody) at dilution 1/700; Secondary antibody: Goat anti rabbit IgG at

Exposure time: 5 seconds

1/8000 dilution;