

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

DDI2 RABBIT PAB

Cat.#: S218584

Product Name: Anti-DDI2 Rabbit Polyclonal Antibody

Synonyms:

UNIPROT ID: Q5TDH0 (Gene Accession - BC006011)

Background: DDI1 and DDI2 are ubiquitin receptor homologs of the Saccharomyces cerevisiae ddi1 protein, which is involved in regulation of the cell cycle and the late secretory pathway. DDI2 is a 399 amino acid protein that contains one ubiquitin-like domain and exists as three isoforms as a result of alternative splicing. The gene encoding DDI2 maps to human chromosome 1, the largest human chromosome which spans about 260 million base pairs and makes up 8% of the human genome. Other notable genes located on chromosome 1 include LMNA, which is associated with the rare aging disease Hutchinson-Gilford progeria, and the MUTYH gene, which is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome.

Immunogen: Fusion protein of human DDI2

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Cell Biology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 218584(DDI2 Antibody) at a dilution of 1/75(Cytoplasm and Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 218584(Anti-DDI2 Antibody) at dilution 1/75.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-2: HL60 and Jurkat cell lysates; Primary antibody: 218584(DDI2 Antibody) at dilution 1/800;

Secondary antibody: HRP-conjugated Goat

anti rabbit IgG at 1/5000 dilution; Exposure time: 30 seconds