

DDX59 RABBIT PAB

Cat.#: S218597

Product Name: Anti-DDX59 Rabbit Polyclonal Antibody

Synonyms: OFD5; ZNHIT5

UNIPROT ID: Q5T1V6 (Gene Accession - BC041801)

Background: DEAD-box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp, are putative RNA helicases implicated in several cellular processes involving modifications of RNA secondary structure and ribosome/spliceosome assembly. Based on their distribution patterns, some members of this family may be involved in embryogenesis, spermatogenesis, and cellular growth and division. DDX59 (DEAD box protein 59), also known as ZNHIT5 (zinc finger HIT domaincontaining protein 5), is a 619 amino acid member of the DEAD box helicase protein family. Like many DEAD box helicase family members, DDX59 contains a Q motif, which controls ATP binding and hydrolysis. Expressed as two isoforms produced by alternative splicing, DDX59 contains one helicase C-terminal domain and one HIT-type zinc finger.

Immunogen: Fusion protein of human DDX59

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 30-150;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG


Purification: Antigen affinity purification

Species Reactivity: Human, Mouse, Rat


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Epigenetics and Nuclear Signaling


Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 218597(DDX59 Antibody) at a dilution of 1/30(Cytoplasm and Nucleus).



In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 218597(Anti-DDX59 Antibody) at dilution 1/30.



Gel: 8%SDS-PAGE, Lysate: 40 μ g;
Lane 1-2: A549 and 293T cell lysates;
Primary antibody: 218597(DDX59 Antibody) at dilution 1/500;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 7 minutes