

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **DKK1 RABBIT PAB**

Cat.#: S210277

Product Name: Anti-DKK1 Rabbit Polyclonal Antibody

Synonyms: SK; DKK-1

UNIPROT ID: 094907 (Gene Accession - BC001539)

**Background:** This gene encodes a member of the dickkopf family of proteins. Members of this family are secreted proteins characterized by two cysteine-rich domains that mediate protein-protein interactions. The encoded protein binds to the LRP6 co-receptor and inhibits beta-catenin-dependent Wnt signaling. This gene plays a role in embryonic development and may be important in bone formation in adults. Elevated expression of this gene has been observed in numerous human cancers and this protein may promote proliferation, invasion and growth in cancer cell lines.

Immunogen: Fusion protein of human DKK1

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

**Purification:** Antigen affinity purification

Species Reactivity: Human, Mouse


**Constituents:** PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

**Research Areas:** Cardiovascular, Immunology, Signal Transduction, Developmental Biology **Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 210277(DKK1 Antibody) at a dilution of 1/50(Secreted).



In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 210277(Anti-DKK1 Antibody) at dilution 1/50.

kDa 95 — 72 — 55 —	
36—	
28—	-
17—	
10 —	

Gel: 10%SDS-PAGE, Lysate: 40 µg; Lane: PC-3 cell lysate; Primary antibody: 210277(DKK1 Antibody) at dilution 1/600; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 40 seconds