

DTYMK RABBIT PAB

Cat.#: S212535

Product Name: Anti-DTYMK Rabbit Polyclonal Antibody

Synonyms: CDC8; TMPK; TYMK; PP3731

UNIPROT ID: P23919 (Gene Accession - BC001827)

Background: DTYMK (deoxythymidylate kinase (thymidylate kinase)), also known as CDC8, TMPK, TYMK or dTMP kinase, is a 212 amino acid protein that belongs to the thymidylate kinase family and is involved in pyrimidine metabolism. Specifically, DTYMK catalyzes the ATP-dependent conversion of dTMP (deoxythymidine monophosphate) to dTDP (deoxythymidine diphosphate), which then functions as one of the four nucleotides in DNA. Via its role in the catalytic creation of dTDP, DTYMK plays an important role in the pathway of DNA synthesis and is thought to be involved in cell cycle progression and cell growth. DTYMK expression levels peak during the S phase (synthesis phase) of the cell cycle, further supporting the role of DTYMK in DNA synthesis.

Immunogen: Fusion protein of human DTYMK

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG


Purification: Antigen affinity purification

Species Reactivity: Human, Mouse


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Epigenetics and Nuclear Signaling


Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 212535(DTYMK Antibody) at a dilution of 1/70(Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 212535(Anti-DTYMK Antibody) at dilution 1/70.



Gel: 8%SDS-PAGE, Lysate: 40 μ g;
 Lane 1-6: Raji, K562, Hela, HepG2, 293T cell and Human fetal liver tissue lysates;
 Primary antibody: 212535(DTYMK Antibody) at dilution 1/550;
 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
 Exposure time: 5 seconds