

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

ELOVL5 RABBIT PAB

Cat.#: S222347

Product Name: Anti-ELOVL5 Rabbit Polyclonal Antibody

Synonyms: HELO1; SCA38; dJ483K16.1

UNIPROT ID: Q9NYP7 (Gene Accession - NP_068586)

Background: This gene belongs to the ELO family. It is highly expressed in the adrenal gland and testis, and encodes a multi-pass membrane protein that is localized in the endoplasmic reticulum. This protein is involved in the elongation of long-chain polyunsaturated fatty acids. Mutations in this gene have been associated with spinocerebellar ataxia-38 (SCA38). Alternatively spliced

transcript variants have been found for this gene.

Immunogen: Synthetic peptide of human ELOVL5

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Cardiovascular, Signal Transduction, Metabolism

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 222347(ELOVL5 Antibody) at a dilution of 1/50(Cytoplasm and Cell membrane).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 222347(Anti-ELOVL5 Antibody) at dilution 1/50.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-7: Mouse fat tissue, Jurkat cell, RAW264.7 cell, Mouse brain tissue, Hela cell, Human testis tissue, Mouse adrenal gland tissue lysates;

Primary antibody: 222347(ELOVL5 Antibody) at

dilution 1/550;

Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 15 seconds