

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

EPSTI1 RABBIT PAB

Cat.#: S217406

Product Name: Anti-EPSTIl Rabbit Polyclonal Antibody

Synonyms: BRESII

UNIPROT ID: Q96J88 (Gene Accession - BC023660)

Background: The protein encoded by this gene has been shown to promote tumor invasion and metastasis in some invasive cancer cells when overexpressed. Expression of this gene has been shown to be upregulated by direct binding of the Kruppel like factor 8 protein to promoter sequences. The translated protein interacts with the amino terminal region of the valosin containing protein gene product, resulting in the nuclear translocation of the nuclear factor kappa B subunit 1 gene product, and activation of target genes. Overexpression of this gene has been observed in some breast cancers and in some individuals with systemic lupus erythematosus (SLE). [provided by RefSeq, Sep 2016]

Immunogen: Fusion protein of human EPSTI

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 100-200;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


Research Areas: Cancer

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 217406(EPSTII Antibody) at a dilution of 1/160(Cytoplasm).

In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the fusion protein and then with 217406(Anti-EPSTII Antibody) at dilution 1/160.

Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: HepG2 cell lysate; Primary antibody: 217406(EPSTII Antibody) at dilution 1/1000; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 1 minute