

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

GFER RABBIT PAB

Cat.#: S216322

Product Name: Anti-GFER Rabbit Polyclonal Antibody

Synonyms: ALR, HPO, HSS, ERV1, HPO1, HPO2, HERV1

UNIPROT ID: P55789 (Gene Accession - BC028348)

Background: The hepatotrophic factor designated augmenter of liver regeneration (ALR) is thought to be one of the factors responsible for the extraordinary regenerative capacity of mammalian liver. It has also been called hepatic regenerative stimulation substance (HSS). The gene resides on chromosome 16 in the interval containing the locus for polycystic kidney disease (PKDI). The putative gene product is 42 similar to the scERVI protein of yeast. The yeast scERVI gene had been found to be essential for oxidative phosphorylation, the maintenance of mitochondrial genomes, and the cell division cycle. The human gene is both the structural and functional homolog of the yeast scERVI gene.

Immunogen: Fusion protein of human GFER

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-200;WB: 500-2000;ELISA: 1000-5000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Signal Transduction, Cell Biology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 216322(GFER Antibody) at a dilution of 1/50(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 216322(Anti-GFER Antibody) at dilution 1/50.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Mouse liver tissue; Primary antibody: 216322(GFER Antibody) at dilution 1/750; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 minute