

## HEXIM1 RABBIT PAB

**Cat.#:** N225448

**Product Name:** Anti-HEXIM1 Rabbit pAb

**Synonyms:** CLP1; EDG1; HIS1; MAQ1

**UNIPROT ID:** O94992

**Background:** Transcriptional regulator which functions as a general RNA polymerase II transcription inhibitor (PubMed:14580347, PubMed:15713661, PubMed:15201869). In cooperation with 7SK snRNA sequesters P-TEFb in a large inactive 7SK snRNP complex preventing RNA polymerase II phosphorylation and subsequent transcriptional elongation (PubMed:12832472, PubMed:14580347, PubMed:15713661, PubMed:15201869). May also regulate NF-kappa-B, ESRI, NR3C1 and CIITA-dependent transcriptional activity (PubMed:15940264, PubMed:15941832, PubMed:17088550). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:28712728).

**Immunogen:** The antiserum was produced against synthesized peptide derived from human HEXIM1. AA range:181-230

**Applications:** WB,IHC-F,IHC-P,ICC/IF,ELISA

**Recommended Dilutions:** WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 ELISA: 1/10000

**Host Species:** Rabbit

**Clonality:** Rabbit Polyclonal

**Clone ID:** -

**MW:** Calculated MW: 41 kDa; Observed MW: 41 kDa

**Isotype:** IgG

**Purification:** Affinity Chromatography

**Species Reactivity:** Human,Mouse,Rat


**Conjugation:** Unconjugated

**Modification:** Unmodified


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**Research Areas:** Epigenetics and Nuclear Signaling


**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing




Immunofluorescence analysis of HEXIM1 in HeLa using HEXIM1 antibody. The picture on the right is blocked using the synthesized peptide.



Western blot analysis of HEXIM1 in rat HEART lysates using HEXIM1 antibody.



Western blot analysis of HEXIM1 in NIH/3T3 lysates using HEXIM1 antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of HEXIM1 in the HeLa lysates using HEXIM1 antibody