

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HTRA2 RABBIT PAB

Cat.#: S221437

Product Name: Anti-HTRA2 Rabbit Polyclonal Antibody

Synonyms: OMI; MGCA8; PARK13; PRSS25

UNIPROT ID: O43464 (Gene Accession - NP_037379)

Background: This gene encodes a serine protease. The protein has been localized in the endoplasmic reticulum and interacts with an alternatively spliced form of mitogen-activated protein kinase 14. The protein has also been localized to the mitochondria with release to the cytosol following apoptotic stimulus. The protein is thought to induce apoptosis by binding the apoptosis inhibitory protein baculoviral IAP repeat-containing 4. Nuclear localization of this protein has also been observed. Alternate splicing of this gene results in multiple transcript variants encoding different isoforms.

Immunogen: Synthetic peptide of human HTRA2

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Cancer, Metabolism, Cell Biology, Neuroscience

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human gastric cancer tissue using 221437(HTRA2 Antibody) at a dilution of 1/50(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with the synthetic peptide and then with 221437(Anti-HTRA2 Antibody) at dilution 1/50.



Gel: 8%SDS-PAGE, Lysate: 40 µg;

Lane: Hela cell lysate;

Primary antibody: 221437(HTRA2 Antibody) at

dilution 1/1000;

Secondary antibody: HRP-conjugated Goat

anti rabbit IgG at 1/5000 dilution;

Exposure time: 3 seconds