

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

IKZF2 RABBIT PAB

Cat.#: S219151

Product Name: Anti-IKZF2 Rabbit Polyclonal Antibody

Synonyms: ANF1A2; HELIOS; ZNF1A2; ZNFN1A2

UNIPROT ID: Q9UKS7 (Gene Accession - BC028936)

Background: This gene encodes a member of the Ikaros family of zinc-finger proteins. Three members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some variants has not been determined.

Immunogen: Fusion protein of human IKZF2

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 100-300;WB: 200-1000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Epigenetics and Nuclear Signaling, Immunology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 219151(IKZF2 Antibody) at a dilution of 1/140(Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 219151(Anti-IKZF2 Antibody) at dilution 1/140.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Human placenta tissue lysate; Primary antibody: 219151(IKZF2 Antibody) at dilution 1/400;

Secondary antibody: HRP-conjugated Goat

anti rabbit IgG at 1/5000 dilution;

Exposure time: 10 minutes