

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **IL9 RABBIT PAB**

Cat.#: S215729

Product Name: Anti-IL9 Rabbit Polyclonal Antibody

Synonyms: P40; HP40; IL-9

UNIPROT ID: P15248 (Gene Accession - NP\_000581)

**Background:** The protein encoded by this gene is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. The gene encoding this cytokine has been identified as a candidate gene for asthma. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the pathogenesis of bronchial hyperresponsiveness.

**Immunogen:** Synthetic peptide of human IL9

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-100;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

**Purification:** Antigen affinity purification

Species Reactivity: Human

**Constituents:** PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


Research Areas: Cancer, Immunology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 215729(IL9 Antibody) at a dilution of 1/30(Secreted).

In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the synthetic peptide and then with 215729(Anti-IL9 Antibody) at dilution 1/30.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane: HL-60 cell lysate; Primary antibody: 215729(IL9 Antibody) at dilution 1/250; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 3 minutes