

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **IL9 RABBIT PAB**

**Cat.#:** S215729

**Product Name:** Anti-IL9 Rabbit Polyclonal Antibody

**Synonyms:** P40; HP40; IL-9

UNIPROT ID: P15248 (Gene Accession - NP\_000581)

**Background:** The protein encoded by this gene is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. The gene encoding this cytokine has been identified as a candidate gene for asthma. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the

pathogenesis of bronchial hyperresponsiveness.

Immunogen: Synthetic peptide of human IL9

**Applications:** ELISA, WB, IHC

**Recommended Dilutions:** IHC: 50-100;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

**Clonality:** Rabbit Polyclonal

**Isotype:** Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification

Species Reactivity: Human

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol

Research Areas: Cancer, Immunology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 215729(IL9 Antibody) at a dilution of 1/30(Secreted).



In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the synthetic peptide and then with 215729 (Anti-IL9 Antibody) at dilution 1/30.



Gel: 12%SDS-PAGE, Lysate: 40 µg;

Lane: HL-60 cell lysate;

Primary antibody: 215729(IL9 Antibody) at

dilution 1/250;

Secondary antibody: HRP-conjugated Goat

anti rabbit IgG at 1/5000 dilution;

Exposure time: 3 minutes