

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

KRT20 RABBIT PAB

Cat.#: S221168

Product Name: Anti-KRT20 Rabbit Polyclonal Antibody

Synonyms: K20; CD20; CK20; CK-20; KRT21

UNIPROT ID: P35900 (Gene Accession - NP_061883)

Background: The protein encoded by this gene is a member of the keratin family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. The type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. This cytokeratin is a major cellular protein of mature enterocytes and goblet cells and is specifically expressed in the gastric and intestinal mucosa. The type I cytokeratin genes are clustered in a region of chromosome 17q12–q21.

Immunogen: Synthetic peptide of human KRT20

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 25-100;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Signal Transduction, Cancer

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human colorectal cancer tissue using 221168 (KRT20 Antibody) at a dilution of 1/20 (Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human colorectal cancer tissue is first treated with the synthetic peptide and then with 221168 (Anti-KRT20 Antibody) at dilution 1/20.



Gel: 12%SDS-PAGE, Lysate: 40 µg;

Lane 1-2: LOVO cell and Mouse small intestines

tissue lysates;

Primary antibody: 221168(KRT20 Antibody) at

dilution 1/600;

Secondary antibody: HRP-conjugated Goat

anti rabbit IgG at 1/5000 dilution;

Exposure time: 2 minutes