

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

LIPG RABBIT PAB

Cat.#: S217402

Product Name: Anti-LIPG Rabbit Polyclonal Antibody
Synonyms: EL; EDL; PRO719
UNIPROT ID: Q9Y5X9 (Gene Accession - BC060825)
Background: The protein encoded by this gene has substantial phospholipase activity and may be involved in lipoprotein metabolism and vascular biology. This protein is designated a member of the

involved in lipoprotein metabolism and vascular biology. This protein is designated a member of the TG lipase family by its sequence and characteristic lid region which provides substrate specificity for enzymes of the TG lipase family.

Immunogen: Fusion protein of human LIPG

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 25-100;WB: 200-1000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human


Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol **Research Areas:** Metabolism, Cardiovascular

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human colorectal cancer tissue using 217402(LIPG Antibody) at a dilution of 1/30(Cytoplasm and Nucleus).




The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using 217402(Anti-LIPG Antibody) at a dilution of 1/30.



In comparision with the IHC on the left, the same paraffin-embedded Human colorectal cancer tissue is first treated with the fusion protein and then with 217402(Anti-LIPG Antibody) at dilution 1/30.



In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with fusion protein and then with D222302(Anti-LIPG Antibody) at dilution 1/30.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Human fetal liver tissue lysate; Primary antibody: 217402(LIPG Antibody) at dilution 1/250; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 2 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010