

## LRRC23 RABBIT PAB

**Cat.#:** S221460

**Product Name:** Anti-LRRC23 Rabbit Polyclonal Antibody

**Synonyms:** LRPB7

**UNIPROT ID:** Q53EV4 (Gene Accession - NP\_964013 )

**Background:** The leucine-rich (LRR) repeat is a 20–30 amino acid motif that forms a hydrophobic  $\beta$ / $\alpha$  horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. The primary function of these motifs is to provide a versatile structural framework to mediate the formation of protein–protein interactions. LRRs are present in a variety of proteins with diverse structure and function, including innate immunity and nervous system development. Several human diseases are associated with mutations in genes encoding LRR-containing proteins. LRRC23 (leucine-rich repeat-containing protein 23), also known as leucine-rich protein B7, is a 343 amino acid protein that contains eight LRR (leucine-rich) repeats and one LRRCT domain. LRRC23 exists as two alternatively spliced isoforms and is encoded by a gene mapping to chromosome 12.

**Immunogen:** Synthetic peptide of human LRRC23

**Applications:** ELISA, IHC

**Recommended Dilutions:** IHC: 25–100; ELISA: 5000–10000

**Host Species:** Rabbit

**Clonality:** Rabbit Polyclonal

**Isotype:** Immunogen-specific rabbit IgG


**Purification:** Antigen affinity purification

**Species Reactivity:** Human


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

**Research Areas:** Cell Biology


**Storage & Shipping:** Store at –20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 221460(LRRC23 Antibody) at a dilution of 1/30(Cell membrane and Nucleus).



In comparison with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the synthetic peptide and then with 221460(Anti-LRRC23 Antibody) at dilution 1/30.



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using 221460(Anti-LRRC23 Antibody) at a dilution of 1/30.



In comparison with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with synthetic peptide and then with D263061(Anti-LRRC23 Antibody) at dilution 1/30.