

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

MAPT RABBIT PAB

Cat.#: S216821

Product Name: Anti-MAPT Rabbit Polyclonal Antibody

Synonyms: TAU; MSTD; PPND; DDPAC; MAPTL; MTBT1; MTBT2; tau-40; FTDP-17; PPP1R103

UNIPROT ID: P10636 (Gene Accession - BC114948)

Background: This gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes comple,x regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia,

cortico-basal degeneration and progressive supranuclear palsy.

Immunogen: Fusion protein of human MAPT

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-200;WB: 1000-5000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Signal Transduction, Neuroscience

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 216821 (MAPT Antibody) at a dilution of 1/75 (Cytoplasm and Cell membrane).



In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the fusion protein and then with 216821(Anti-MAPT Antibody) at dilution 1/75.



Gel: 6%SDS-PAGE, Lysate: 40 µg; Lane: Mouse fetal brain tissue lysate; Primary antibody: 216821(MAPT Antibody) at dilution 1/1000;

Secondary antibody: HRP-conjugated Goat

anti rabbit IgG at 1/5000 dilution;

Exposure time: 5 seconds