

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

MRPS18C RABBIT PAB

Cat.#: S218090

Product Name: Anti-MRPS18C Rabbit Polyclonal Antibody

Synonyms: CGI-134; S18mt-c; MRPS18-1; mrps18-c; MRP-S18-1; MRP-S18-c

UNIPROT ID: Q9Y3D5 (Gene Accession - BC005186)

Background: Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology.

Immunogen: Full length fusion protein

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 25-100;WB: 1000-5000;ELISA: 2000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Epigenetics and Nuclear Signaling

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 218090(MRPS18C Antibody) at a dilution of 1/25(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 218090(Anti-MRPS18C Antibody) at dilution 1/25.



Gel: 12%SDS-PAGE, Lysate: 40 µg;

Lane 1-4: 231, Hela, K562 cell and Human

bladder carcinoma tissue;

Primary antibody: 218090 (MRPS18C Antibody)

at dilution 1/800;

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution;

Exposure time: 5 seconds