

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## NDUFA8 RABBIT PAB

Cat.#: S220736

Product Name: Anti-NDUFA8 Rabbit Polyclonal Antibody

Synonyms: PGIV; CI-19KD; CI-PGIV

UNIPROT ID: P51970 (Gene Accession - NP\_055037)

**Background:** The protein encoded by this gene belongs to the complex I 19 kDA subunit family. Mammalian complex I is composed of 45 different subunits. This protein has NADH dehydrogenase activity and oxidoreductase activity. It plays an important role in transfering electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone. **Immunogen:** Synthetic peptide of human NDUFA8

Applications: ELISA, WB, IHC

**Recommended Dilutions:** IHC: 50-200;WB: 500-2000;ELISA: 2000-5000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human


**Constituents:** PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol **Research Areas:** Metabolism, Cancer, Cell Biology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human prostate cancer tissue using 220736(NDUFA8 Antibody) at a dilution of 1/40(Cytoplasm).



paraffin-embedded Human brain tissue using 220736(Anti-NDUFA8 Antibody) at a dilution of 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with the synthetic peptide and then with 220736(Anti-NDUFA8 Antibody) at dilution 1/40.



The image on the left is immunohistochemistry of In comparision with the IHC on the left, the same paraffin-embedded Human brain tissue is first treated with synthetic peptide and then with D261941(Anti-NDUFA8 Antibody) at dilution 1/40.



Gel: 10%SDS-PAGE, Lysate: 40 µg; Lane: 293T cells; Primary antibody: 220736(NDUFA8 Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 minut



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010