

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

PDIA4 RABBIT PAB

Cat.#: S218903

Product Name: Anti-PDIA4 Rabbit Polyclonal Antibody

Synonyms: ERP70; ERP72; ERp-72

UNIPROT ID: P13667 (Gene Accession - BC000425)

Background: This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, three catalytically active thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence. This protein, when bound to cyclophilin B, enhances the rate of immunoglobulin G intermolecular disulfide bonding and antibody assembly.

Immunogen: Fusion protein of human PDIA4

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 40-200;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Signal Transduction

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 218903(PDIA4 Antibody) at a dilution of 1/40(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the fusion protein and then with 218903(Anti-PDIA4 Antibody) at dilution 1/40.





The image on the left is immunohistochemistry of In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue paraffin-embedded Human thyroid cancer tissue using 218903(Anti-PDIA4 Antibody) at a dilution of is first treated with fusion protein and then with 1/40.

D225471(Anti-PDIA4 Antibody) at dilution 1/40.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-2: RAW264.7 and NIH/3T3 cell lysates; Primary antibody: 218903(PDIA4 Antibody) at dilution 1/250; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 second



Product Description

Pioneering GTPase and Oncogene Product Development since 2010