

PIK3CD RABBIT PAB

Cat.#: S220808

Product Name: Anti-PIK3CD Rabbit Polyclonal Antibody

Synonyms: APDS; PI3K; IMD14; p110D; IMD14A; IMD14B; ROCHIS; PI10DELTA

UNIPROT ID: O00329 (Gene Accession - NP_005017)

Background: Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I PI3K found primarily in leukocytes. Like other class I PI3Ks (p110-alpha p110-beta, and p110-gamma), the encoded protein binds p85 adapter proteins and GTP-bound RAS. However, unlike the other class I PI3Ks, this protein phosphorylates itself, not p85 protein.

Immunogen: Synthetic peptide of human PIK3CD

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-200;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG


Purification: Antigen affinity purification

Species Reactivity: Human, Mouse


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Signal Transduction, Immunology


Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 220808(PIK3CD Antibody) at a dilution of 1/50(Cytoplasm).




In comparison with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 220808(Anti-PIK3CD Antibody) at dilution 1/50.



The image on the left is immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using 220808(Anti-PIK3CD Antibody) at a dilution of 1/50.



In comparison with the IHC on the left, the same paraffin-embedded Human colorectal cancer tissue is first treated with synthetic peptide and then with D262049(Anti-PIK3CD Antibody) at dilution 1/50.



Gel: 6%SDS-PAGE, Lysate: 40 µg;
Lane: 293T cell lysate;
Primary antibody: 220808(PIK3CD Antibody) at dilution 1/800;
Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;
Exposure time: 2 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
