

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## PSMA6 RABBIT PAB

Cat.#: S219283

**Product Name:** Anti-PSMA6 Rabbit Polyclonal Antibody

Synonyms: IOTA; p27K; PROS27

UNIPROT ID: P60900 (Gene Accession - BC002979)

**Background:** The proteasome is a multicatalytic proteinase complex with a highly ordered ringshaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase TIA family, that is a 20S core alpha subunit. Multiple transcript variants encoding several different isoforms have been found for this gene. A pseudogene has been identified on the Y chromosome.

Immunogen: Fusion protein of human PSMA6

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 100-300;WB: 1000-5000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human, Mouse, Rat

**Constituents:** PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


Research Areas: Cell Biology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 219283(PSMA6 Antibody) at a dilution of 1/90(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the fusion protein and then with 219283(Anti-PSMA6 Antibody) at dilution 1/90.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-5: A375, HL60 cell, Mouse skeletal muscle tissue, Jurkat, 293T cell lysates; Primary antibody: 219283(PSMA6 Antibody) at dilution 1/1000; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 90 seconds