

PYRUVATE DEHYDROGENASE E1 ALPHA RABBIT MAB

Cat.#: N262790

Product Name: Anti-Pyruvate Dehydrogenase E1 alpha Rabbit Monoclonal Antibody

Synonyms: mitochondrial; ODP_A_HUMAN; PDH; PDHA; PDHA1; PDHCE1A; PDHE1 A type I; PDHE1-A type I; PHE1A; Pyruvate Dehydrogenase (lipoamide) alpha 1; Pyruvate dehydrogenase complex; E1 alpha polypeptide 1; Pyruvate Dehydrogenase E1 alpha; Pyruvate dehydrogenase E1 component subunit alpha; Pyruvate dehydrogenase E1 component subunit alpha; somatic form; mitochondrial; somatic form.

UNIPROT ID: P08559

Background: The PDH complex is composed of multiple copies of three enzymatic components: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and lipoamide dehydrogenase (E3). Catalyzes the removal of CO₂ from pyruvate. Mutations in the α subunits of pyruvate dehydrogenase (E1) lead to congenital defects that are usually associated with lactic acidosis, neurodegeneration and early death.

Immunogen: A synthetic peptide of human Pyruvate Dehydrogenase E1-alpha subunit

Applications: WB,ICC/IF,IP

Recommended Dilutions: WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20

Host Species: Rabbit

Clonality: Rabbit Monoclonal

Clone ID: R06-1J4

MW: Calculated MW: 43 kDa; Observed MW: 43 kDa

Isotype: IgG

Purification: Affinity Purified

Species Reactivity: Human,Mouse,Rat


Conjugation: Unconjugated

Modification: Unmodified


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

Research Areas: Signal Transduction

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Immunocytochemistry analysis of Pyruvate Dehydrogenase E1alpha subunit (green) in Hela using Pyruvate Dehydrogenase E1alpha subunit antibody, and DAPI (blue)



Western blot analysis of Pyruvate Dehydrogenase E1alpha subunit in C6, 3T3 lysates using Pyruvate Dehydrogenase E1 alpha antibody.