

## RAD54L2 RABBIT PAB

**Cat.#:** S216760

**Product Name:** Anti-RAD54L2 Rabbit Polyclonal Antibody

**Synonyms:** ARIP4; HSPC325; SRISNF2L

**UNIPROT ID:** Q9Y4B4 (Gene Accession - BC024298 )

**Background:** DNA helicase that modulates androgen receptor (AR)-dependent transactivation in a promoter-dependent manner. Not able to remodel mononucleosomes in vitro (By similarity).

**Immunogen:** Fusion protein of human RAD54L2

**Applications:** ELISA, WB, IHC

**Recommended Dilutions:** IHC: 50-200;WB: 200-1000;ELISA: 2000-5000

**Host Species:** Rabbit

**Clonality:** Rabbit Polyclonal

**Isotype:** Immunogen-specific rabbit IgG


**Purification:** Antigen affinity purification

**Species Reactivity:** Human, Mouse


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

**Research Areas:** Signal Transduction, Epigenetics and Nuclear Signaling


**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 216760(RAD54L2 Antibody) at a dilution of 1/60(Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 216760(Anti-RAD54L2 Antibody) at dilution 1/60.



Gel: 6%SDS-PAGE, Lysate: 40  $\mu$ g;  
Lane 1-4: Hela cells, Raji cells, 293T cells, PC3 cells;  
Primary antibody: 216760(RAD54L2 Antibody) at dilution 1/400;  
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;  
Exposure time: 5 minutes