

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## SEMA6D RABBIT PAB

Cat.#: S220893

Product Name: Anti-SEMA6D Rabbit Polyclonal Antibody

Synonyms:

**UNIPROT ID:** Q8NFY4 (Gene Accession - NP\_705872)

**Background:** Semaphorins are a large family, including both secreted and membrane associated proteins, many of which have been implicated as inhibitors or chemorepellents in axon pathfinding, fasciculation and branching, and target selection. All semaphorins possess a semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in the N-terminal extracellular portion. Additional sequence motifs C-terminal to the semaphorin domain allow classification into distinct subfamilies. Results demonstrate that transmembrane semaphorins, like the secreted ones, can act as repulsive axon guidance cues. This gene encodes a class 6 vertebrate transmembrane semaphorin that demonstrates alternative splicing. Several transcript variants have been identified and expression of the distinct encoded isoforms is thought to be regulated in a tissue- and development-dependent manner.

Immunogen: Synthetic peptide of human SEMA6D

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-100;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human, Mouse

**Constituents:** PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


Research Areas: Cancer, Neuroscience

**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 220893(SEMA6D Antibody) at a dilution of 1/30(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 220893(Anti-SEMA6D Antibody) at dilution 1/30.

kDa	
250—	
130—	-
95—	
72—	
55—	
36 —	

Gel: 6%SDS-PAGE, Lysate: 40 µg; Lane: LOVO cell lysate; Primary antibody: 220893(SEMA6D Antibody) at dilution 1/700; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 5 minutes