

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

SLC22A6 RABBIT PAB

Cat.#: S220156

Product Name: Anti-SLC22A6 Rabbit Polyclonal Antibody

Synonyms: OAT1, PAHT, HOAT1, ROAT1

UNIPROT ID: Q4U2R8 (Gene Accession - NP_004781)

Background: The protein encoded by this gene is involved in the sodium-dependent transport and excretion of organic anions, some of which are potentially toxic. The encoded protein is an integral membrane protein and may be localized to the basolateral membrane. Four transcript variants encoding four different isoforms have been found for this gene. Functions as organic anion exchanger when the uptake of one molecule of organic anion is coupled with an efflux of one molecule of endogenous dicarboxylic acid (glutarate, ketoglutarate, etc). Mediates the sodium-independent uptake of 2,3-dimercapto-1-propanesulfonic acid (DMPS) By similarity.

Immunogen: Synthetic peptide of human SLC22A6

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-200;WB: 1000-5000;ELISA: 2000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification

Species Reactivity: Human, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Metabolism

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 220156(SLC22A6 Antibody) at a dilution of 1/50(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the synthetic peptide and then with 220156(Anti-SLC22A6 Antibody) at dilution 1/50.



Gel: 10%SDS-PAGE, Lysate: 40 µg; Lane: Human fetal kidney tissue;

Primary antibody: 220156(SLC22A6 Antibody)

at dilution 1/900;

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution;

Exposure time: 3 minutes