

STK3/STK4 RABBIT PAB

Cat.#: S220426

Product Name: Anti-STK3/STK4 Rabbit Polyclonal Antibody

Synonyms: KRS1; MST2/KRS2; MST1; YSK3; TIIAC

UNIPROT ID: Q13043 (Gene Accession - NP_006273)

Background: Ste20 (Ste20) is a serine/threonine kinase in *Saccharomyces cerevisiae* that is involved in relaying signals from G protein-coupled receptors to cyto-solic MAP kinase cascades. Mammalian protein kinases that display sequence similarity to Ste20 are divided into two groups, the PAK subfamily and the GCK subfamily. The mammalian Ste20-like kinases (MST kinases), also known as Krs proteins, are members of the GCK subfamily. Ksr-1 (MST-2) and Ksr-2 (MST-1) are both direct substrates of caspase-3 that accelerate caspase-3 activation.

Immunogen: Synthetic peptide of human STK3/STK4

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 100-300;WB: 500-2000;ELISA: 2000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG


Purification: Antigen affinity purification

Species Reactivity: Human, Mouse


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Signal Transduction, Cancer


Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human Lymphoma tissue using 220426(STK3/STK4 Antibody) at a dilution of 1/70(Cytoplasm).




In comparison with the IHC on the left, the same paraffin-embedded Human Lymphoma tissue is first treated with the synthetic peptide and then with 220426(Anti-STK3/STK4 Antibody) at dilution 1/70.



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using 220426(Anti-STK3/STK4 Antibody) at a dilution of 1/70.



In comparison with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with synthetic peptide and then with D261514(Anti-STK3/STK4 Antibody) at dilution 1/70.



Gel: 6%SDS-PAGE, Lysate: 40 µg;
 Lane 1-5: PC3 cells, hela cells, human bladder carcinoma tissue, A172 cells, hepG2 cells;
 Primary antibody: 220426(STK3/STK4 Antibody) at dilution 1/1000;
 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
 Exposure time: 10 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
