


TRAF3 RABBIT PAB**Cat.#:** S220010**Product Name:** Anti-TRAF3 Rabbit Polyclonal Antibody**Synonyms:** CAPI, LAPI, CAP-1, CRAF1, IIAE5, CD40bp**UNIPROT ID:** Q13114 (Gene Accession - NP_003291.2)

Background: The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from, members of the TNF receptor (TNFR) superfamily. This protein participates in the signal transduction of CD40, a TNFR family member important for the activation of the immune response. This protein is found to be a critical component of the lymphotoxin-beta receptor (LTbetaR) signaling complex which induces NF-kappaB activation and cell death initiated by LTbeta ligation. Epstein-Barr virus encoded latent infection membrane protein-1 (LMP1) can interact with this and several other members of the TRAF family, which may be essential for the oncogenic effects of LMP1. Several alternatively spliced transcript variants encoding three distinct isoforms have been reported.


Immunogen: Synthetic peptide of human TRAF3**Applications:** ELISA, WB, IHC**Recommended Dilutions:** IHC: 25-100;WB: 200-1000;ELISA: 1000-5000**Host Species:** Rabbit**Clonality:** Rabbit Polyclonal**Isotype:** Immunogen-specific rabbit IgG**Purification:** Antigen affinity purification**Species Reactivity:** Human, Mouse**Constituents:** PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol**Research Areas:** Signal Transduction, Cancer, Immunology**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human tonsil tissue using 220010(TRAF3 Antibody) at a dilution of 1/25(Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with the synthetic peptide and then with 220010(Anti-TRAF3 Antibody) at dilution 1/25.



Gel: 8+10+12%SDS-PAGE, Lysate: 60 µg;
Lane: Mouse kidney tissue;
Primary antibody: 220010(TRAF3 Antibody) at dilution 1/300;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 40 seconds