

TRANSCRIPTION FACTOR AP 2 GAMMA RABBIT MAB

Cat.#: N261851

Product Name: Anti-Transcription Factor AP 2 gamma Rabbit Monoclonal Antibody

Synonyms: ERF1; TFAP2G; hAP-2g; AP2-GAMMA

UNIPROT ID: Q92754

Background: The protein encoded by this gene is a sequence-specific DNA-binding transcription factor involved in the activation of several developmental genes. The encoded protein can act as either a homodimer or heterodimer with other family members and is induced during retinoic acid-mediated differentiation. It plays a role in the development of the eyes, face, body wall, limbs, and neural tube.

Immunogen: Recombinant protein of human AP2 gamma

Applications: WB, ICC/IF, IP

Recommended Dilutions: WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20

Host Species: Rabbit

Clonality: Rabbit Monoclonal

Clone ID: R03-812

MW: Calculated MW: 49 kDa; Observed MW: 49 kDa

Isotype: IgG

Purification: Affinity Purified

Species Reactivity: Human


Conjugation: Unconjugated

Modification: Unmodified


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

Research Areas: Cell Biology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Immunocytochemistry analysis of TFAP2C/AP2 gamma (green) in MCF-7 using TFAP2C/AP2 gamma antibody, and DAPI (blue)



Western blot analysis of TFAP2C/AP2 gamma in Jurkat, Hela lysates using Transcription Factor AP 2 gamma antibody.