

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

TXNDC12 RABBIT PAB

Cat.#: S219224

Product Name: Anti-TXNDC12 Rabbit Polyclonal Antibody

Synonyms: AGI; AGRI; ERP16; ERP18; ERP19; TLP19; hAG-1; PDIA16; hTLP19

UNIPROT ID: 095881 (Gene Accession - BC008913)

Background: This gene encodes a member of the thioredoxin superfamily. Members of this family are characterized by a conserved active motif called the thioredoxin fold that catalyzes disulfide bond formation and isomerization. This protein localizes to the endoplasmic reticulum and has a single atypical active motif. The encoded protein is mainly involved in catalyzing native disulfide bond formation and displays activity similar to protein-disulfide isomerases. This protein may play a role in defense against endoplasmic reticulum stress. Alternate splicing results in both coding and non-coding variants.

Immunogen: Fusion protein of human TXNDC12

Applications: ELISA, IHC

Recommended Dilutions: IHC: 50-200; ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human, Mouse, Rat

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


Research Areas: Signal Transduction

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human colorectal cancer tissue using 219224(TXNDC12 Antibody) at a dilution of 1/70(Cytoplasm). In comparision with the IHC on the left, the same paraffin-embedded Human colorectal cancer tissue is first treated with the fusion protein and then with 219224(Anti-TXNDC12 Antibody) at dilution 1/70.



The image on the left is immunohistochemistry of paraffinembedded Human liver cancer tissue using 219224(Anti-TXNDC12 Antibody) at a dilution of 1/70.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with fusion protein and then with D226086(Anti-TXNDC12 Antibody) at dilution 1/70.