

## UBIQUITIN RABBIT PAB

**Cat.#:** N226046

**Product Name:** Anti-Ubiquitin Rabbit pAb

**Synonyms:** FLJ25987; MGC8385; ubiquitin B; Ubiquitin; UBCEP1; UBCEP2; RPS27A

**UNIPROT ID:** P0CG47

**Background:** Plays an important role in the ubiquitin-proteasome pathway. Ubiquitin can be covalently linked to many cellular proteins by the ubiquitination process, which targets proteins for degradation by the 26S proteasome. Three components are involved in the target protein-ubiquitin conjugation process. Ubiquitin is first activated by forming a thiolester complex with the activation component E1; the activated ubiquitin is subsequently transferred to the ubiquitin-carrier protein E2, then from E2 to ubiquitin ligase E3 for final delivery to the epsilon-NH<sub>2</sub> of the target protein lysine residue.

**Immunogen:** A synthetic peptide of human Ubiquitin

**Applications:** WB,IHC-P,FC

**Recommended Dilutions:** WB: 1/500-1/1000 IHC: 1/50-1/100 FC: 1/50-1/100

**Host Species:** Rabbit

**Clonality:** Rabbit Polyclonal

**Clone ID:** -

**MW:** Calculated MW: 26 kDa; Observed MW: 8 kDa

**Isotype:** IgG

**Purification:** Affinity Purified

**Species Reactivity:** Human


**Conjugation:** Unconjugated

**Modification:** Unmodified


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**Research Areas:** Neuroscience

**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin-embedded Human Cholangiocarcinoma using Ubiquitin antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

 Western blot analysis of Ubiquitin in 293 lysates using Ubiquitin antibody.