

## UBL7 RABBIT PAB

**Cat.#:** S217970

**Product Name:** Anti-UBL7 Rabbit Polyclonal Antibody

**Synonyms:** TCBA1; BMSC-UbP

**UNIPROT ID:** Q96S82 (Gene Accession - BC007913 )

**Background:** Ubiquitin is an important cellular signal that targets proteins for degradation or regulates their functions. UBL7 (ORF name SB132) encodes ubiquitin-like protein7 also named bone marrow stromal cell ubiquitin-like protein (BMSC-UbP) capable of binding ubiquitin. BMSC-UbP protein is derived from bone marrow stromal cells containing a ubiquitin-associated (UBA) domain at the C terminus that has been implicated in linking cellular processes and the ubiquitin system. BMSC-UbP mRNA is widely expressed in human multiple tissues and various tumor cell lines and is decreased in BMSC stimulated with phorbol myristate acetate (PMA) suggesting a role in regulation of BMSC function.

**Immunogen:** Fusion protein of human UBL7

**Applications:** ELISA, WB, IHC

**Recommended Dilutions:** IHC: 100-200;WB: 500-2000;ELISA: 5000-10000

**Host Species:** Rabbit

**Clonality:** Rabbit Polyclonal

**Isotype:** Immunogen-specific rabbit IgG


**Purification:** Antigen affinity purification

**Species Reactivity:** Human, Mouse


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

**Research Areas:** Cell Biology


**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 217970(UBL7 Antibody) at a dilution of 1/100(Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the fusion protein and then with 217970(Anti-UBL7 Antibody) at dilution 1/100.



Gel: 8%SDS-PAGE, Lysate: 40 µg;  
Lane 1-2: Mouse testis tissue, Hela cell lysates;  
Primary antibody: 217970(UBL7 Antibody) at dilution 1/700;  
Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;  
Exposure time: 1 minute