

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

WNK2 RABBIT PAB

Cat.#: S221121

Product Name: Anti-WNK2 Rabbit Polyclonal Antibody **Synonyms:** PRKWNK2; NY-CO-43; SDCCAG43; P/OKcl.13 **UNIPROT ID:** Q9Y3S1 (Gene Accession - NP_001269323)

Background: The protein encoded by this gene is a cytoplasmic serine-threonine kinase that belongs to the protein kinase superfamily. The protein plays an important role in the regulation of electrolyte homeostasis, cell signaling survival, and proliferation. Alternative splicing results in multiple transcript

variants.

Immunogen: Synthetic peptide of human WNK2

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 25-100;WB: 200-1000;ELISA: 1000-2000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification **Species Reactivity:** Human, Mouse


Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

Research Areas: Signal Transduction, Epigenetics and Nuclear Signaling **Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 221121(WNK2 Antibody) at a dilution of 1/35(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the synthetic peptide and then with 221121(Anti-WNK2 Antibody) at dilution 1/35.



The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using 221121(Anti-WNK2 Antibody) at a dilution of 1/35.



In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with synthetic peptide and then with D262500(Anti-WNK2 Antibody) at dilution 1/35.



Gel: 6%SDS-PAGE, Lysate: 40 µg; Lane 1-2: A431 cells, MCF7 cells;

Primary antibody: 221121(WNK2 Antibody) at

dilution 1/200;

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution;

Exposure time: 3 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010