

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

XPR1 RABBIT PAB

Cat.#: S213296

Product Name: Anti-XPR1 Rabbit Polyclonal Antibody

Synonyms: X3; SYG1

UNIPROT ID: Q9UBH6 (Gene Accession - NP_004727)

Background: Xenotropic mouse leukemia viruses (X-MLVs) are broadly infectious for mammals except most of the classical strains of laboratory mice. These gammaretroviruses rely on the XPR1 receptor for entry, and the unique resistance of laboratory mice is due to two mutations in different putative XPR1extracellular loops. Cells from avian species differ in susceptibility to X-MLVs, and 2 replacement mutations in the virus-resistant chicken XPR1 distinguish it from the more permissive duck and quail receptors.

Immunogen: Synthetic peptide of human XPR1

Applications: ELISA, IHC

Recommended Dilutions: IHC: 50-200; ELISA: 1000-5000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG

Purification: Antigen affinity purification

Species Reactivity: Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


Research Areas: Signal Transduction

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human ovarian cancer tissue using 213296(XPR1 Antibody) at a dilution of 1/45(Cytoplasm).



The image on the left is immunohistochemistry of paraffinembedded Human gastric cancer tissue using 213296(Anti-XPR1 Antibody) at a dilution peptide and then with D152188(Anti-XPR1 of 1/45.



In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with the synthetic peptide and then with 213296(Anti-XPRI Antibody) at dilution 1/45.



In comparision with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with synthetic Antibody) at dilution 1/45.