ATM PAB

ATM Polyclonal Antibody

Cat. #: 21276
Gene Symbol: ATM
Description: Anti-ATM Rabbit Polyclonal Antibody
Background: Ataxia telangiectasia mutated (ATM) is a serine/threonine protein kinase that is recruited and activated by DNA double-strand breaks. It phosphorylates several key proteins that initiate activation of the DNA damage checkpoint, leading to cell cycle arrest, DNA repair or apoptosis. Activity of ATM protein is under tight control, and mutation of ATM can cause disease such as Ataxia telangiectasia (AT) and cancers.
Immunogen: Purified ATM protein, human origin.
Applications: ELISA, WB, IHC
Recommended Dilutions:
ELISA: 1:1000-1:5000
WB: 1:100-1:1000
IHC: 1:50-1:200
Concentration: 0.2 mg/ml
Host Species: Rabbit
Format: Liquid
Clonality: Polyclonal
Isotype:
Purity: Purified from serum
Preservative: No
Constituents: PBS (without Mg^{2+} and Ca^{2+}), pH 7.4, 150 mM NaCl, 50% glycerol
Species Reactivity: Recognizes ATM of vertebrates.
Storage Conditions: Store at -20°C. Avoid repeated freezing and thawing

Western blot:

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS



Western blot analysis of recombinant ATM protein. Purified His-tagged ATM protein was blotted with Anti-ATM rabbit polyclonal antibody (Cat. #21276).

Immunohistochemistry:



Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue with Anti-ATM polyclonal antibody (Cat. #21276). Tissue samples were fixed with formaldehyde and blocked with 1% serum for 15 min at 37 °C. Antigen retrieval was by heat mediation in citrate buffer (pH6). Samples were then incubated with primary antibody (1:50 dilution) overnight at 4°C. A HRP-conjugated Goat anti-mouse IgG (1:50 dilution) was used as secondary antibody.

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS