

# **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

### **B-CATENIN(E53K)**

### **β-Catenin(E53K)**

**Cat. #:** 26167

Gene Symbol: CTNNB1, CTNNB

**Description:** Anti-β-catenin(E53K) Mouse Monoclonal Antibody

**Background:** Beta-catenin (or  $\beta$ -catenin) is a protein that in humans is encoded by the CTNNB1 gene.  $\beta$ -catenin is a subunit of the cadherin protein complex and also acts as an intracellular signal transducer in the Wnt signaling pathway. Deregulation of beta-catenin signaling is an important event in the genesis of a number of malignancies, such as colon cancer, melanoma, hepatocellular carcinoma, ovarian cancer, endometrial cancer, medulloblastoma pilomatricomas, and prostate cancer.

**Immunogen:** A synthetic peptide from the internal region of  $\beta$ -catenin which includes the mutation of E53K, human origin.

Applications: ELISA, WB, IF, IHC

**Recommended Dilutions:** 

ELISA: 1:1000-1:5000 WB: 1:500-1:2000 IHC: 1:50-1:100

Concentration: 1 mg/ml Host Species: Mouse

Format: Liquid

**Clonality:** Monoclonal

Isotype: IgG

Purity: Purified from ascites

**Preservative:** No

Constituents: PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 50%

glycerol

**Species Reactivity:** Recognizes  $\beta$ -catenin(E53K), but not wild type  $\beta$ -catenin

protein from vertebrates.

Storage Conditions: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

#### **Western blot:**

# β-catenin protein WT E53K




1 2

### WB: Anti-β-catenin (E53K) mAb

Western blot analysis of recombinant β-catenin(E53K) and wild type proteins. Purified His-tagged β-catenin(E53K) protein (amino acids 1-76, lane 2) and corresponding wild type protein (lane 1) were blotted with Anti-β-catenin(E53K) mouse monoclonal antibody (Cat. #26167).

#### Immunofluorescence:



Immunofluorescence of cells expressing  $\beta$ -catenin proteins with anti- $\beta$ -catenin(E53K) antibody. HEK293T cells were transfected with pCDNA3-GFP- $\beta$ -catenin (WT) plasmid (left column) or pCDNA3-GFP- $\beta$ -catenin(E53K) plasmid (right column), then fixed and stained with Anti- $\beta$ -catenin (E53K) monoclonal antibody (Cat. #26167).