

CD162 (DMC283) IGG1 CHIMERIC MAB

Cat.#: 28191

Product Name: Anti-CD162(DMC283) IgG1 Chimeric Monoclonal Antibody

Synonyms: PSGL-1;PSGL1;CD162;SELPLG;Selectin P ligand

Description: Anti-CD162 antibody(DMC283) IgG1 Chimeric Monoclonal Antibody

Background: This gene encodes a glycoprotein that functions as a high affinity counter-receptor for the cell adhesion molecules P-; E- and L-selectin expressed on myeloid cells and stimulated T lymphocytes. As such; this protein plays a critical role in leukocyte trafficking during inflammation by tethering of leukocytes to activated platelets or endothelia expressing selectins. This protein requires two post-translational modifications; tyrosine sulfation and the addition of the sialyl Lewis x tetrasaccharide (sLex) to its O-linked glycans; for its high-affinity binding activity. Aberrant expression of this gene and polymorphisms in this gene are associated with defects in the innate and adaptive immune response. Alternate splicing results in multiple transcript variants.

Applications: Flow Cyt

Recommended Dilutions: Flow Cyt 1:100

Host Species: Rabbit

Isotype: Rabbit:Human Fc chimeric IgG1

Purification: Purified from cell culture supernatant by affinity chromatography

Species Reactivity: Human CD162

Constituents: Lyophilized from sterile PBS, pH 7.4. 5 % – 8% trehalose is added as protectants before lyophilization.

Storage & Shipping: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. Flow cytometry analysis with Anti-CD162 (DMC283) on Expi293 cells transfected with human CD162 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).