

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## CD164 (DMC476) IGG1 CHIMERIC MAB

Cat.#: 28252

Product Name: Anti-CD164(DMC476) IgG1 Chimeric Monoclonal Antibody

**Synonyms:** LMOR; M-OR-1; MOP; MOR; MOR1; OPRM

**Description:** Anti-CD164 antibody(DMC476) IgG1 Chimeric Monoclonal

**Antibody** 

**Background:** This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as betaendorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse; such as nicotine; cocaine; and alcohol via its modulation of the dopamine system. The NM\_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains. [provided by RefSeq; Oct 2013]

**Applications:** Flow Cyt

Recommended Dilutions: Flow Cyt 1:100

Host Species: Rabbit

Isotype: Rabbit:Human Fc chimeric IgG1

Purification: Purified from cell culture supernatant by affinity

chromatography

Species Reactivity: Human CD164

Constituents: Lyophilized from sterile PBS, pH 7.4. 5 % – 8% trehalose is

added as protectants before lyophilization.

**Storage & Shipping:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010




Figure 1. CD164 protein is highly expressed on the surface of Expi293 cell membrane. Flow cytometry analysis with Anti-CD164 (DMC476) on Expi293 cells transfected with human CD164 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram), and Isotype antibody on Expi293 transfected with irrelevant protein (Orange histogram).