

## DNAM-1 (DM96) RABBIT MAB

**Cat.#:** 28434

**Product Name:** Anti-DNAM-1(DM96) Rabbit Monoclonal Antibody

**Synonyms:** DNAM1; CD226; PTA1

**Description:** Anti-DNAM-1 antibody(DM96) Rabbit Monoclonal Antibody

**Background:** This gene encodes a glycoprotein expressed on the surface of NK cells; platelets; monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. The protein mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Alternative splicing results in multiple transcript variants.

**Applications:** ELISA; Flow Cyt

**Recommended Dilutions:** ELISA 1:5000-10000; Flow Cyt 1:100

**Host Species:** Rabbit

**Isotype:** Rabbit IgG

**Purification:** Purified from cell culture supernatant by affinity chromatography

**Species Reactivity:** Human DNAM1

**Constituents:** Lyophilized from sterile PBS, pH 7.4. 5% - 8% trehalose is added as protectants before lyophilization.

**Storage & Shipping:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human DNAM-1 protein, mFc-His tagged protein 11176 can bind Rabbit anti-DNAM-1 monoclonal antibody (clone: DM96) in a linear range of 0.64-80 ng/ml.




Figure 2. Expi 293 cell line transfected with irrelevant protein (A) and human DNAM-1 (B) were surface stained with Rabbit anti-DNAM-1 monoclonal antibody 1 µg/ml (clone: DM96) followed by PE-conjugated anti-rabbit IgG secondary antibody.