

ERBB2 PAB

ErbB2 Polyclonal Antibody

Cat. #: 21202

Gene Symbol: ERBB2/HER2/Neu

Description: Anti-ErbB2 Rabbit Polyclonal Antibody

Background: The ErbB family is composed of four plasma membrane-bound receptor tyrosine kinases. Amplification or over-expression of the ERBB2 gene occurs in approximately 30% of breast cancers. HER2 is encoded by ERBB2, a known proto-oncogene located at the long arm of human chromosome 17(17q21-q22). HER2 proteins have been shown to form clusters in cell membranes that may play a role in tumorigenesis.

Immunogen: A synthetic peptide from the central of ErbB2 protein.

Applications: ELISA, WB, IHC

Recommended Dilutions:

ELISA: 1:500-1:5000

WB: 1:200-1:2000

IHC: 1:20-1:200

Concentration: 0.2 mg/ml

Host Species: Rabbit

Format: Liquid

Clonality: Polyclonal

Isotype: IgG

Purity: Purified from serum

Preservative: No


Constituents: PBS (without Mg^{2+} and Ca^{2+}), pH 7.4, 150 mM NaCl, 50% glycerol

Species Reactivity: Recognizes ErbB2 of vertebrates.

Storage Conditions: Store at $-20^{\circ}C$. Avoid repeated freezing and thawing.


Western blot:

ERBB2 protein



WB: Anti-ERBB2 pAb

Western blot analysis of ErbB2 protein. Purified full length His-tagged ErbB2 protein (Cat. #10208) was blotted with Anti-ErbB2 rabbit polyclonal antibody (Cat. #21202).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue with Anti-ErbB2 polyclonal antibody (Cat. #21202). Tissue samples were fixed with formaldehyde and blocked with 1% serum for 15 min at 37 °C. Antigen retrieval was by heat mediation in citrate buffer (pH6). Samples were then incubated with primary antibody (1:50 dilution) overnight at 4 °C. A HRP-conjugated Goat anti-Rabbit IgG (1:50) was used as secondary antibody.